

Ordinary vs Double Schubert polynomials

Shizuo KAJI (Yamaguchi University)

Thm

There is an explicit relationship between ordinary and double Schubert polynomials

Torus equivariant cohomology of flag varieties

$G \supset T \curvearrowright G/B$: multiplication

$H_T^*(G/B; \mathbb{R})$ is freely generated by the Schubert classes \mathfrak{S}_w ($w \in W$: Weyl gp.) over $H^*(BT; \mathbb{R}) = \mathbb{R}[t_1, \dots, t_n]$

$$H_T^*(G/B; \mathbb{R}) \cong \bigoplus_{w \in W} \mathbb{R}[t_1, \dots, t_n] \langle \mathfrak{S}_w \rangle$$

(1) free module

On the other hand, as an algebra over $H^*(BT; \mathbb{R})$,

$$H_T^*(G/B; \mathbb{R}) \cong \frac{\mathbb{R}[t_1, \dots, t_n, x_1, \dots, x_n]}{(f(x) - f(t))}$$

(f : runs all the W -invariants)

(2) coinvariant ring of W

Two different presentations

Schubert polynomials

The *double Schubert polynomials* (of 2-sets of variables) are polynomial representatives of Schubert classes in (2)

$$\mathfrak{S}_w(t; x) \in \frac{\mathbb{R}[t_1, \dots, t_n, x_1, \dots, x_n]}{(f(x) - f(t))}$$

Similarly, the *ordinary Schubert polynomials* (of 1-set of variables) are for the ordinary cohomology.

$$\sigma_w(x) \in \frac{\mathbb{R}[x_1, \dots, x_n]}{(f(x))} \cong H^*(G/B; \mathbb{R})$$

A construction of σ_w is given uniformly for all Lie-types in Bernstein-Gelfand-Gelfand '73

Schubert polynomial relates the two presentations

Double \Rightarrow Ordinary

There are two ways to obtain the ordinary Schubert polynomials from the double ones:

(i) $\sigma_w(x) = \mathfrak{S}_w(0; x)$

(ii) $\sigma_w(x) = \frac{1}{|W|} \sum_{v \in W} \mathfrak{S}_{w^{-1}v}(-x; v^{-1}(-x))$

This is not surprising since G/B is a *GKM-manifold*, and hence the equivariant cohomology contains all the information of the ordinary cohomology

Equivariant cohomology is a finer invariant

Ordinary \Rightarrow Double

We can also go the other way.

Let the *partition of $w \in W$ into i parts* be a set

$$P_i(w) = \{(w_1, w_2, \dots, w_i) \in W^i \mid w_1 \cdot w_2 \cdots w_i = w, l(w_k) > 0 \forall k, l(w_1) + \dots + l(w_i) = l(w)\}$$

Then

$$\mathfrak{S}_w(t; x) = \sum_{i=1}^{l(w)} \sum_{(w_1, w_2, \dots, w_i) \in P_i(w)} (-1)^{i-1} \sigma_{w_1}(t) \sigma_{w_2}(t) \cdots \sigma_{w_{i-1}}(t) (\sigma_{w_i}(t) - \sigma_{w_i}(x))$$

In fact, ordinary cohomology recovers equivariant one

Key machinery : Characterization of Schubert classes by the localization to the fixed points and the divided difference operators