

# Low rank cohomology of the classifying spaces of gauge groups over 3-manifolds

By

Shizuo KAJI \*

## Abstract

The purpose of this paper is to calculate the cohomology of the function space  $\text{Map}(M, BG)$  for degree less than or equal to 3, where  $G$  is a simply connected compact Lie group and  $M$  is a closed orientable 3-manifold. The calculation enables us to obtain a simple proof and an improvement of the result [4, Theorem 1.2].

## §1. Introduction

Let  $G$  be a simply connected compact Lie group and  $M$  a closed orientable 3-manifold. Since  $BG$  is 3-connected, any principal  $G$ -bundles over  $M$  are trivial. Then we call the gauge group of the trivial  $G$ -bundle over  $M$  the gauge group over  $M$  and denote it by  $\mathcal{G}$ . It is well-known that

$$B\mathcal{G} \simeq \text{Map}(M, BG)$$

([2]). The cohomology of  $B\mathcal{G}$  in low dimensions is considered in [4] by making use of the Eilenberg-Moore spectral sequence.

**Theorem 1.1** [4, Theorem 1.2]. *Suppose that  $\text{Tor}_{\mathbb{Z}}(\mathbb{Z}/2, R) = 0$ . Let  $G$  be a simply-connected compact Lie group such that the integral cohomology of  $BG$  is torsion free and let  $M$  be a closed orientable 3-manifold. We denote  $H^i(\text{Map}(M, BG); R)$  by  $H^i$ . Then there exists a short exact sequence*

$$0 \rightarrow H_1(M; R)^{\oplus r} \oplus R^{\oplus s} \oplus H^1 \otimes H^2 \xrightarrow{\alpha} H^3 \rightarrow (R/2R)^{\oplus r} \rightarrow 0,$$

where  $\alpha|_{H^1 \otimes H^2}$  is the cup product,  $r = \text{rank}H^4(BG)$  and  $s = \text{rank}H^6(BG)$ . Moreover  $H^1$  is a free  $R$ -module for any  $R$ , and  $H^2$  is also free if  $R$  is a PID.

---

Communicated by K. Saito. Received December 17, 2004. Revised March 25, 2005.  
2000 Mathematics Subject Classification(s): 55R40, 54C35

\*Department of Mathematics, Kyoto University, Kyoto 606-8502, Japan.

The purpose of this paper is to refine Theorem 1.1 and to give a simple proof. We determine the integral cohomology of  $H^i(\text{Map}(M, BG))(i \leq 3)$  without the assumption that  $H^*(BG)$  is torsion free. It is known that  $G$  is the direct product of simply connected compact simple Lie groups ([6]). Then we reduce Theorem 1.1 to the case that  $G$  is a simply connected compact simple Lie group and obtain

**Theorem 1.2.** *Let  $G$  be a simply connected compact simple Lie group and  $M$  a closed orientable 3-manifold. We denote  $H^i(\text{Map}(M, BG))$  by  $H^i$ . Then we have*

$$H^i \cong \begin{cases} \mathbb{Z} & i = 0 \\ H^1(\Omega^2 G) & i = 1 \\ H_2(M) & i = 2 \\ H^1(\Omega^2 G) \otimes H_2(M) \oplus H_1(M) \oplus H^3(\Omega^2 G) & i = 3. \end{cases}$$

Moreover, the cup product  $H^1 \otimes H^2 \rightarrow H^3$  maps  $H^1 \otimes H^2$  isomorphically onto the direct summand  $H^1(\Omega^2 G) \otimes H_2(M) \subset H^3$ .

*Remark.* Let  $G$  be a simply connected compact simple Lie group. We here describe the integral cohomology  $H^*(\Omega^2 G)$  for  $* \leq 3$ .

Table 1.  $H^*(\Omega^2 G)$

$H^i(\Omega^2 G)$	type of $G$		
	$A_l(l \geq 2)$	$C_l(l \geq 1)$	otherwise
$i = 1$	$\mathbb{Z}$	$\mathbb{Z}$	$\mathbb{Z}$
$i = 2$	0	0	0
$i = 3$	$\mathbb{Z}$	$\mathbb{Z}/2$	0

*Remark.* Since the inclusion of the 1-skeleton  $\bigvee^{g-1} S^1 \rightarrow \overbrace{\mathbb{R}P^2 \# \dots \# \mathbb{R}P^2}^g$  induces an isomorphism on mod  $p$  cohomology for each odd prime  $p$ , Theorem 1.1 in [4] is easily shown without the assumption that  $H^*(G)$  is  $p$ -torsion free by [5, Proposition 4.2] and [3, Ch. VI, Proposition 7.1].

## §2. Approximation of $G$ by infinite loop spaces

Let  $G$  be a simply connected compact Lie group. In this section we approximate  $G$  by an infinite loop space in low dimensions.

It is known that  $G$  is the direct product of simply connected compact simple Lie groups. Since simply connected compact simple Lie groups are classified by their Lie algebras as  $A_l, B_l, C_l, D_l (l \geq 1), E_l (l = 6, 7, 8), F_4, G_2$ , we give an approximation to each type.

**Proposition 2.1.** *Let  $G$  be a simply connected compact simple Lie group. Then there exist an infinite loop space  $\mathbf{B}$  and a 7-equivalence  $t : BG \rightarrow \mathbf{B}$ .*

*Proof.* First we note that there are the following correspondence in low ranks ([6]):

$$A_1 = B_1 = C_1, B_2 = C_2, D_2 = A_1 \times A_1, D_3 = A_3.$$

Therefore we have only to consider the types  $A_l (l \geq 2), B_l (l \geq 3), C_l (l \geq 1), D_l (l \geq 4), E_l (l = 6, 7, 8), F_4$  and  $G_2$ . In the case where  $G$  is of type  $A_l (l \geq 2)$ , there exists a 7-equivalence  $BG \rightarrow BSU$  induced by the inclusion  $G \rightarrow SU$ . Similarly in the case where  $G$  is of type  $C_l (l \geq 1)$ , there exists a 7-equivalence  $BG \rightarrow BSp$ . For  $G$  otherwise, we have  $\pi_i(G) = 0$  ( $i = 1, 2, 4, 5$ ) ([6]). Then a representative of a generator of  $H^4(BG; \mathbb{Z}) \cong \mathbb{Z}$  is a 7-equivalence  $BG \rightarrow K(\mathbb{Z}, 4)$ .  $\square$

**Corollary 2.1.** *Let  $M$  be a 3-dimensional complex. Then we have  $H^i(\text{Map}(M, \mathbf{B})) \cong H^i(\text{Map}(M, BG))$  ( $i \leq 3$ ), where  $\mathbf{B}$  is as in Proposition 2.1.*

*Proof.* Let  $\text{Map}_*(X, Y)$  denote the space of basepoint preserving maps from  $X$  to  $Y$ , where  $X, Y$  are based spaces.

We consider the following commutative diagram

$$\begin{array}{ccc} \pi_k(\text{Map}_*(M, BG)) & \xrightarrow{\pi_*(\text{Map}_*(Id, t))} & \pi_k(\text{Map}_*(M, \mathbf{B})) \\ \cong \downarrow & & \cong \downarrow \\ [S^k \wedge M, BG] & \xrightarrow{t_*} & [S^k \wedge M, \mathbf{B}]. \end{array}$$

Since the second row is an isomorphism for  $k \leq 3$  and a surjection for  $k = 4$  by J.H.C. Whitehead theorem, we see that  $\text{Map}_*(Id, t) : \text{Map}_*(M, BG) \rightarrow \text{Map}_*(M, \mathbf{B})$  is a 4-equivalence. Consider the following commutative diagram

of evaluation fibrations

$$\begin{array}{ccccc}
\mathrm{Map}_*(M, BG) & \longrightarrow & \mathrm{Map}(M, BG) & \longrightarrow & BG \\
\mathrm{Map}_*(Id, t) \downarrow & & \mathrm{Map}(Id, t) \downarrow & & t \downarrow \\
\mathrm{Map}_*(M, \mathbf{B}) & \longrightarrow & \mathrm{Map}(M, \mathbf{B}) & \longrightarrow & \mathbf{B}
\end{array}$$

Since  $\mathrm{Map}_*(Id, t)$  is a 4-equivalence and  $t$  is a 7-equivalence, it follows that  $\mathrm{Map}(Id, t) : \mathrm{Map}(M, BG) \rightarrow \mathrm{Map}(M, \mathbf{B})$  is a 4-equivalence.  $\square$

### §3. Proof of Theorem 1.2

Let  $G$  be a simply connected compact simple Lie group. By Proposition 2.1 there exists an infinite loop space  $\mathbf{B}$  and a 7-equivalence  $t : BG \rightarrow \mathbf{B}$ . Since  $\mathbf{B}$  is a homotopy group, we have a homotopy equivalence

$$\begin{aligned}
\mathrm{Map}(M, \mathbf{B}) &\simeq \mathrm{Map}_*(M, \mathbf{B}) \times \mathbf{B} \\
f &\mapsto (f \cdot f(*)^{-1}, f(*)),
\end{aligned}$$

where  $*$  denotes the basepoint of  $M$ . Since the infinite loop space  $\mathbf{B}$  is 3-connected, the inclusion map  $\mathrm{Map}_*(M, B) \rightarrow \mathrm{Map}(M, B)$  induces the isomorphism on homology for degree less than or equal to 3. Then we compute  $H^*(\mathrm{Map}_*(M, \mathbf{B}))$  to determine  $H^*(\mathrm{Map}(M, \mathbf{B}))$ .

**Proposition 3.1.** *We have*

$$H^k(\mathrm{Map}_*(M, \mathbf{B})) \cong \bigoplus_{i+j=k} H^i(\mathrm{Map}_*(M^2, \mathbf{B})) \otimes H^j(\Omega^3 \mathbf{B})$$

for  $k \leq 3$ , where  $M^2$  is the 2-skeleton of  $M$ .

*Proof.* Since a closed orientable 3-manifold is parallelizable, the top cell of  $M$  is split off stably ([1]). Actually by Freudenthal suspension theorem the top cell of  $M$  is split off after double suspension. Then we have

$$\begin{aligned}
\mathrm{Map}_*(M, \mathbf{B}) &\simeq \mathrm{Map}_*(M, \Omega^2 B^2 \mathbf{B}) \\
&\simeq \mathrm{Map}_*(\Sigma^2 M, B^2 \mathbf{B}) \\
&\simeq \mathrm{Map}_*(\Sigma^2 M^2 \vee S^5, B^2 \mathbf{B}) \\
&\simeq \mathrm{Map}_*(M^2 \vee S^3, \mathbf{B}) \\
&\simeq \mathrm{Map}_*(M^2, \mathbf{B}) \times \Omega^3 \mathbf{B}.
\end{aligned}$$

Since  $H^k(\Omega^3 \mathbf{B})$  ( $k < 3$ ) is either 0 or  $\mathbb{Z}$ , the proof is completed by Künneth Theorem.  $\square$

To compute  $H^i(\text{Map}_*(M^2, \mathbf{B}))$  ( $i \leq 3$ ) we need the following technical lemma. Let  $X, Y, Z$  be based spaces and  $f : X \rightarrow Y$  be a based map. We denote by  $f^\#$  the induced map  $\text{Map}_*(f, Id) : \text{Map}_*(Y, Z) \rightarrow \text{Map}_*(X, Z)$ .

**Lemma 3.1.** *Let  $X$  be a based space such that there is a  $(p + q + 1)$ -equivalence  $g : X \rightarrow K(\mathbb{Z}, p + q)$  and let  $f : \bigvee^l S^p \rightarrow \bigvee^m S^p$  be a based map. Suppose  $(f^\#)^* : H^q(\Pi^l \Omega^p X) \rightarrow H^q(\Pi^m \Omega^p X)$  is represented by a matrix  $A$  for a certain basis. Then  $f_* : H_p(\bigvee^l S^p) \rightarrow H_p(\bigvee^m S^p)$  is also represented by  $A$  for a suitable basis.*

*Proof.* Since  $g_* : [S^{p+q}, X] \rightarrow [S^{p+q}, K(\mathbb{Z}, p + q)]$  is an isomorphism, we have the following commutative diagram

$$\begin{array}{ccccccc} H_q(\Pi^m \Omega^p X) & \xrightarrow[\text{hur}]{\cong} & \pi_q(\Pi^m \Omega^p X) & \xlongequal{\quad} & [\bigvee^m S^{p+q}, X] & \xrightarrow{\cong} & H^{p+q}(\bigvee^m S^{p+q}) \\ (f^\#)_* \downarrow & & (f^\#)_* \downarrow & & \downarrow & & (\Sigma^q f)^* \downarrow \\ H_q(\Pi^l \Omega^p X) & \xrightarrow[\text{hur}]{\cong} & \pi_q(\Pi^l \Omega^p X) & \xlongequal{\quad} & [\bigvee^l S^{p+q}, X] & \xrightarrow{\cong} & H^{p+q}(\bigvee^l S^{p+q}), \end{array}$$

where  $\text{hur}$  is the Hurewicz homomorphism. Since  $\Omega^p X \rightarrow K(\mathbb{Z}, q)$  is a  $(q + 1)$ -equivalence, the proof is completed by taking the dual.  $\square$

$$\mathbf{Proposition\ 3.2.} \quad H^i(\text{Map}_*(M^2, \mathbf{B})) \cong \begin{cases} 0 & i = 1 \\ H_2(M) & i = 2 \\ H_1(M) & i = 3 \end{cases}$$

*Proof.* We have the following cofibration sequence  $\bigvee^l S^1 \xrightarrow{f} \bigvee^m S^1 \xrightarrow{i} M^2 \rightarrow \bigvee^l S^2$ , where  $f$  is the attaching map of 2-cells of  $M$  and  $i$  is the inclusion. Then we have the fibration

$$\Pi^l \Omega^2 \mathbf{B} \rightarrow \text{Map}_*(M^2, \mathbf{B}) \xrightarrow{i^\#} \Pi^m \Omega \mathbf{B}.$$

We consider the Leray-Serre spectral sequence  $(E_r, d_r)$  of the fibration above. Since  $E_2^{p,q} \cong H^p(\Pi^m \Omega \mathbf{B}) \otimes H^q(\Pi^l \Omega^2 \mathbf{B})$ ,  $\mathbf{B}$  is 3-connected and  $H^4(\mathbf{B}) \cong \mathbb{Z}$ , the non-trivial differential  $d_r : E_r^{p,q} \rightarrow E_r^{p+r, q-r+1}$  ( $p + q \leq 4$ ) occurs only when  $r = 3$  and  $(p, q) = (0, 2)$ . Then we obtain  $H^1(\text{Map}_*(M^2, \mathbf{B})) = 0$ . Next we determine  $d_3 : E_3^{0,2} \rightarrow E_3^{3,0}$  to compute  $H^i(\text{Map}_*(M^2, \mathbf{B}))$  ( $i = 2, 3$ ). We

consider the commutative diagram

$$\begin{array}{ccccccc}
\mathbb{V}^l S^1 & \xrightarrow{f} & \mathbb{V}^m S^1 & \xrightarrow{i} & M^2 & \longrightarrow & \mathbb{V}^l S^2 \\
f \downarrow & & 1 \downarrow & & \downarrow & & \Sigma f \downarrow \\
\mathbb{V}^m S^1 & \xrightarrow{1} & \mathbb{V}^m S^1 & \xrightarrow{j} & \mathbb{V}^m D^2 & \longrightarrow & \mathbb{V}^m S^2,
\end{array}$$

where  $j$  is the inclusion. Applying  $\text{Map}_*(\cdot, \mathbf{B})$  to above, we have the following commutative diagram

$$\begin{array}{ccccc}
\Pi^m \Omega \mathbf{B} & \xleftarrow{i^\#} & \text{Map}_*(M^2, \mathbf{B}) & \xleftarrow{\quad} & \Pi^l \Omega^2 \mathbf{B} \\
1 \uparrow & & \uparrow & & (\Sigma f)^\# \uparrow \\
\Pi^m \Omega \mathbf{B} & \xleftarrow{j^\#} & \Pi^m P(\Omega \mathbf{B}) & \xleftarrow{\quad} & \Pi^m \Omega^2 \mathbf{B},
\end{array}$$

where the second row is the product of the path space fibrations of  $\Omega \mathbf{B}$ . Comparing the Leray-Serre spectral sequence of fibrations above, we obtain  $d_3 = \tau(\Sigma f)^\#_* : E_3^{0,2} \rightarrow E_3^{3,0}$ , where  $\tau : H^2(\Pi^m \Omega^2 \mathbf{B}) \xrightarrow{\cong} H^3(\Pi^m \Omega \mathbf{B})$  is the transgression.

Let  $A$  be a matrix which represents  $((\Sigma f)^\#)^* : H^2(\Pi^l \Omega^2 \mathbf{B}) \rightarrow H^2(\Pi^m \Omega^2 \mathbf{B})$ . By Lemma 3.1,  $(\Sigma f)_* : H_2(\mathbb{V}^l S^2) \rightarrow H_2(\mathbb{V}^m S^2)$  is represented by  $A$  and so is  $f_* : H_1(\mathbb{V}^l S^1) \rightarrow H_1(\mathbb{V}^m S^1)$ . Then we have the exact sequence

$$0 \rightarrow H_2(M^2) \rightarrow H_1(\mathbb{V}^l S^1) \xrightarrow{A} H_1(\mathbb{V}^m S^1) \rightarrow H_1(M^2) \rightarrow 0.$$

Since  $H_i(M^2) \cong H_i(M)$  ( $i \leq 2$ ), we have

$$\begin{aligned}
H^2(\text{Map}_*(M^2, \mathbf{B})) &\cong \text{Ker}\{d_3 : E_3^{0,2} \rightarrow E_3^{3,0}\} \cong \text{Ker} A \cong H_2(M), \\
H^3(\text{Map}_*(M^2, \mathbf{B})) &\cong \text{Coker}\{d_3 : E_3^{0,2} \rightarrow E_3^{3,0}\} \cong \text{Coker} A \cong H_1(M).
\end{aligned}$$

□

*Proof of Theorem 1.2.* By Corollary 2.1, Proposition 3.1 and Proposition 3.2, Theorem 1.2 is proved. □

#### §4. Acknowledgment

I would like to thank Akira Kono and Daisuke Kishimoto for their helpful conversation and useful comments.

### References

- [1] Atiyah, M., Thom complexes, *Proc. London Math. Soc. (3)*, **11** (1961), 291–310.
- [2] Atiyah, M., Bott, R., The Yang-Mills equations over Riemann surfaces, *Philos. Trans. Roy. Soc. London Ser. A.*, **308** (1982), 523–615.
- [3] Bousfield, A. K., Kan, D. M., Homotopy limits, completions and localizations, *Lecture Notes in Math.*, **304**, Springer-Verlag, Berlin and New York (1972).
- [4] Kuribayashi, K., Eilenberg-Moore spectral sequence calculation of function space cohomology, *manuscripta math.*, **114** (2004), 305–325.
- [5] May, J., Fibrewise localization and completion, *Trans. Amer. Math. Soc.*, **258** no. 1 (1980), 127–146.
- [6] Mimura, M., Toda, H., Topology of Lie groups. I, II, *Translated from the 1978 Japanese edition by the authors, Translations of Mathematical Monographs*, **91**, American Mathematical Society, Providence, RI (1991).